4.0 SITE EVALUATIONS AND RANKINGS

4.1 Use of Evaluation Criteria

While there are numerous potential evaluation parameters and many possible levels of weighting within each, there are no standard methods by which all wetland sites can be evaluated objectively, resulting in a single list of preservation priorities that is scientifically based and defensible. The many different important resource and site factors involved dictate a more subjective approach in assessing priorities for wetland preservation. To the extent that objective criteria can be integrated into an evaluation procedure (either to rate potential preserve and mitigation sites, or to determine the appropriateness of development), there should be site-by-site consideration of all factors, although sometimes certain factors may be weighted differently for different subregions or resource situations. Further, there are few enough sites left that support the listed species and/or constitute relatively natural situations, that a rigorous evaluation and ranking effort does not seem warranted or particularly necessary to establish priorities.

4.2 Site Rankings

No rigorous ranking or single list of sites have been compiled as part of this study. Rather, several lists of sites (see Appendix D) supporting endangered species have been prepared, in generally descending order of size and significance. It should be recognized that there are numerous different resource situations that warrant protection, and it is difficult to compare (in terms of the need for preservation) small sites with sensitive species against large areas of pristine habitat without such species, or urban fragments with little waterfowl value against rural properties with good wildlife value. Examples of situations that may warrant protection, but might not rank high based on a generic evaluation scheme include solitary occurrences of unique species assemblages, special soil situations, geographic outlyers, especially large or unique populations, and areas of exemplary habitat even without sensitive species currently in evidence.

General ranking of sites can be made, including overall ratings (largely proportional to sensitive species occurrences and site size), but may be more useful if subdivided into categories that could include (1) each of the endangered species, (2) soil and geographic uniqueness and representation, (3) the degree of demonstrable threat, (4) site size and extent of habitat to be saved, (5) sites with high restoration potential, and finally once individual site evaluations have been made, (6) availability and cost.

4.3 Priorities for Preservation

While there are numerous factors that could be analyzed for possible rankings of the known sites, there are relatively few large, high quality sites left in the study area (C. Patterson, unpubl. field data; B. Guggolz and M. Waaland, pers. comm.). Because of the limited number of sites known to still support sizable populations of sensitive species and/or significant acreage of relatively undisturbed vernal pool habitat, the priorities for preservation include virtually all sites with viable populations of sensitive species, generally in descending order based on acreage (from larger to smaller sites). Site size is a major consideration, with small sites (generally less than about 5 acres) being much more vulnerable to outside influences and adjacent land use disturbances. Such small sites, especially if isolated or in urban areas, present difficult management and defensibility issues, and generally offer low potential to preserve, restore, or conserve wetland wildlife values. With regard to botanical resources and the less mobile aquatic animal species, however, even small sites can represent viable situations. Even these represent management problems, and the long term viability of such preserves is unknown.

Availability (i.e., willing sellers) is a major issue in this regard, and acquisition efforts will likely need to address sites with willing sellers before others. With a severely limited universe of sites containing known high quality resources, however, theoretical acquisition priorities should be determined on the basis of biological site quality, plus the degree of imminent threat, size and location, and defensibility/manageability.

In spite of the numerous parameters identified as being relevant to site quality (including the extreme variability within some), one level of preservation priority must address the different sensitive species at issue. Specifically, separate priorities should be identified for each species of major concern, in addition to sites with species ensembles or extensive wetland habitat. For example, while there may be a number of good potential preserve sites that contain one particular species, preservation priorities should recognize the need for preservation of other species occurrences, even if they occur on smaller, more degraded properties. Similarly, the full geographic distribution of any given sensitive species should be considered in prioritizing acquisitions. Consideration should be given to the range of soil types and hydrological systems, with all significant situations represented in the eventual preserve system, rather than simply acquiring several large parcels in one particular area or in one (more common) situation.

The ultimate preserve system needs to include the full range of soils and habitat conditions, good geographic representation, and as many large undisturbed properties as possible. Once a regional analysis of the remaining sites is made and the various location, quality, and site characteristics are sorted into categories that need to be represented, the superimposing of availability will likely result in a relatively short list of priority sites. Finally, even with all of the data collection, site evaluations and comparisons that are possible, the ultimate acquisition of properties for inclusion into the regional wetland preserve system will likely hinge upon availability, acquisition funding, and professional judgment.

4.4 Priorities for Restoration

There is an abundance of land in the study area that is suitable for restoration. The greatest potential is land that has suitable clay soil and relatively undisturbed topography and hydrology. Grazing lands, irrigated (but ungraded) hayfields, and lightly cropped properties represent the best opportunities for restoration in that simple removal of the current disruptive land uses would probably result in considerable natural recovery of the wetlands on these lands. Some active restoration efforts may be necessary to remove well established weeds and reestablish native plant species, but with natural drainage patterns and suitable soils still intact, such efforts could be highly successful.

Lands which have been disked, cultivated, graded or filled may also have potential for restoration, but closer investigation would be warranted to determine the potential degree of recovery possible before embarking on major restoration efforts. If soils and/or hydrology have been altered, restoration may be much more difficult if not impossible. Minimally altered topography can often be returned to suitable terrain for wetlands, and even much of the region's cultivated Clear Lake clay could be reconfigured to foster wetland formation.

A distinction should be made in relation to restoration in that the reestablishment of true vernal pool communities may not be possible in some situations, while other seasonal wetland habitats and values may be attainable. Each type of wetland has its own inherent values, and restoration efforts should recognize that some sites offer potential for some types and not others. Restoration efforts, therefore, need to clearly identify the goals and types of habitats to be established, and sites being considered for restoration need to be examined with the various levels of potential in mind. Where vernal pools per se are involved, more rigorous site investigations may be warranted to determine suitability and to direct the design work. Greater detail may be needed in designing a new drainage/habitat reconfiguration that truly represents the pre-disturbance grades and avoids situating the restored pools on previously non-wetland ground.

In response to several recent development projects, wetland habitat <u>creation</u> has also been attempted on the Plain. Such habitat creation occurs on land where minimal wetlands now exist and where historic wetlands may not be evident. If suitable soils and terrain are available, such efforts can result in the establishment of valuable wetland communities. This type of effort generally requires significant soil removal and careful grading to attain a shallow layer of loam over clay, configured in low gradient swales and basins. Details of habitat creation efforts are not presented here, but such efforts constitute a potentially useful means of retaining net wetland acreage in the study area as well as providing suitable new habitat for native plants and animals.

References and Literature Cited

- Barbour, M. G. and J. Major (eds.). 1977. Terrestrial vegetation of California. Wiley Interscience; New York, London, Sidney, Toronto.
- Bauder, T. B. 1986. San Diego vernal pools: recent and projected losses, their condition, and threats to their existence, 1979-1990. Prepared for the Endangered Plant Program, Calif. Dept. of Fish and Game, Sacramento, CA
- Belk, D. 1975. Key to the Anostraca (fairy shrimp) of North America. The Southwestern Naturalist 20 (1): 91-103.
- Brown, C. and S. K. Jain. 1977. Status report on Limnanthes vinculans. Unpubl. report compiled for the California Native Plant Society, Sacramento, CA.
- But, P. 1977. Pleuropogon hooverianus, a status report. Unpubl. status report to the California Native Plant Society, Sacramento.
- Calif. Natural Diversity Data Base. 1988. Unpubl. file data available at Calif. Dept. of Fish and Game, Sacramento, CA.
- Calif. Natural Diversity Data Base. 1989. Data reports produced by the Calif. Natural Diversity Data Base for the Santa Rosa region, including sensitive plant species mapping and occurrence reports. June 19, 1985; Calif. Dept. Fish and Game, Non-Game Heritage Section, Sacramento, CA.
- Calif. Natural Diversity Data Base. 1994. Endangered and Threatened Animals of California.

 List of state and federal Endangered and Threatened animals of California. April, 1994. California Dept. of Fish and Game, Natural Heritage Division, Natural Diversity Data Base, Sacramento, CA.
- California Dept. of Fish and Game. 1988. Information leaflet, California's fully protected birds, mammals, reptiles, amphibians, and fish. Leaflet/memo clarifying state policy, March 1988. CDFG, Sacramento, CA
- California Dept. of Fish and Game. 1989. 1988 annual report on the status of California's state listed Threatened and Endangered plants and animals. Unpubl. report from the State of Calif., The Resources Agency, Sacramento, CA
- California Dept. of Fish and Game. 1994. Endangered, Threatened, and Rare plants of California.

 List of plants, April, 1994. Calif. Dept. of Fish and Game, Natural Heritage Division, Endangered Plant Program, Sacramento, CA.
- California Dept. of Fish and Game. (no date) 1991 annual report on the status of California state listed threatened and endangered animals and plants. Unpubl. report from the State of Calif., The Resources Agency, Sacramento, CA
- California Native Plant Society. Ongoing Collection of rare plant maps for California.

 On file at the Calif. Natural Diversity Data Base, Dept. of Fish and Game, Sacramento, CA.
- Cardwell, G. T. 1958. Geology and ground water in the Santa Rosa and Petaluma Valley areas, Sonoma County, California. U.S. Geological Survey water supply paper 1427; USGS, San Francisco.
- Cowardin, L. M. et al. 1979. Classification of wetlands and deepwater habitats of the United States. FWS/OBS-79/31 (Service Admin. Manual, 30 AM 10.0). U.S. Fish and Wildlife Service.
- de Mars, J, M Johnson, T Lipshutz, G Madrigal, M Roberts. 1977. Laguna de Santa Rosa environmental analysis and management plan. Senior project in Environmental Studies and Planning, Sonoma State University, Rohnert Park
- Eakins, D. G. 1993. Comparison of the aquatic invertebrate fauna of natural and artificial vernal pools in the Santa Rosa Plain, Sonoma County, California. Unpubl. report to the California Department of Fish and Game, Yountville.

- EIP Associates. 1993. Southwest Area Plan Environmental Impact Report. Draft EIR for the City of Santa Rosa's Southwest Area Plan.
- Environmental Science Associates, Inc. 1991. Environmental Impact Report for the San Miguel Estates No.s 1, 2 and 3; TMD Brown subdivision; NWSR 2-87 Annexation. Prepared for the City of Santa Rosa; SCH No. 91043035
- Fahselt, D. 1988. The dangers of transplantation as a conservation technique. Natural Areas Journal 8 (4) 238-244.
- Hanes, T. 1991. Letter to Ralph Osterling Consultants; Re: Hydrological assessment of the Santa Rosa Golf Course Cramer property. September 3, 1991.
- Harrison, N. 1991. A flora of selected sites of the Laguna de Santa Rosa.

 Unpubl. plant checklist and discussion of vernal pools and vernally wet meadows.
- Heady, H. F. 1977. Valley grassland. Chapter 14 In: Barbour and Major (eds.). 1977. Terrestrial vegetation of California. Wiley-Interscience.
- Heiser, C, B., Jr. 1947. A new species of Blennosperma from California. Madrono Vol. 9 103-104.
- Helley, E. J., K. R. Lajoie, W. E. Spangle, M. L. Blair. 1979. Flatland deposits of the San Francisco Bay region, California; their geology and engineering properties and their importance to comprehensive planning. U.S. Geological Survey Professional Paper 943. USGS, San Francisco, 88 pp.
- Higgins, C. G. 1952. Lower course of the Russian River, California.

 Univ. Calif. Publications in Geological Sciences, Vol. 29, No. 5, pp. 181-264. Univ. Calif.
- Holland, R. F. 1978. The geographic and edaphic distribution of vernal pools in the Great Central Valley, California. Calif. Native Plant Society Special Publication No. 4
- Holland, R. F. 1986. Preliminary descriptions of the terrestrial natural communities of California. Unpubl. paper prep. for the Dept. of Fish and Game, Nongame Heritage Program, The Resources Agency, Sacramento, CA
- Holland, R. F. and S. K. Jain. 1977. Vernal pools. Chapter 15 In: Barbour and Major (eds).1977. Terrestrial vegetation of California. Wiley Interscience, New York, London, Sydney, Toronto.
- Holsinger, J. R. 1972. The freshwater amphipod crustaceans (Gammaridae) of North America.

 Biota of freshwater ecosystems, Identification Manual 5; U.S. Environmental Protection Agency. 89 pp.
- Jain, S. (ed.). 1976. Vernal pools, their ecology and conservation. Institute of Ecology Publication Number 9; Univ. Calif. Davis
- Jain, S. and J. P. Moyle (eds.). 1981. Vernal pools and intermittent streams. Institute of Ecology Publication Number 28; Univ. Calif. Davis.
- Jepson, W. L. 1925. A manual of the flowering plants of California. Univ. Calif. Press. Berkeley. 1238 pp.
- LeBaron, G. and J. Mitchell. 1993. Santa Rosa, a 20th Century town. Historia Ltd.

- Long, M. M., M. Friley, D. Densmore, J. De Weese. 1992. Wetland losses within northern California from projects authorized under Nationwide permit No. 26. Unpubl. report prepared by the U.S. Fish and Wildlife Service.
- Lovell, B. L. 1980. Limnanthes vinculans, an account of known field research in Sonoma County. Unpubl. report to the California Natural Diversity Data Base; Sacramento.
- LSA Associates. 1991. Southwest Santa Rosa Master Environmental Assessment. Prepared for the City of Santa Rosa. April, 1991.
- Mason, H. L. 1975. A flora of the marshes of California. Univ. of California Press, Berkeley, Los Angeles, CA.
- McCafferty, W. P. 1981. Aquatic entomology. The fisherman's and ecologist's illustrated guide to insects and their relatives. Science Books International, Boston, Mass.. 448 pp.
- Miller, V. C. (ed.) 1972. Soil survey: Sonoma County California.Prep. by USDA-Soil Conservation Service and US Forest Service in cooperation with Univ. Calif. Agricultural Experiment Station. May, 1972.
- Munz, P. A. 1968. Supplement to a California flora. Univ. of Calif. Press., Berkeley, Los Angeles, CA.
- Munz, P. A., and D. D. Keck. 1959. A California flora. Univ. of Calif. Press. Berkeley, Los Angeles, London.
- National Technical Committee for Hydric Soils. 1987. Hydric soils of the United States. USDA-SCS National Bulletin No. 430-5-9. USDA-SCS, Washington, DC
- Nejedly, J. 1977. Native Plant Protection: Senate Bill 308, Chapter 1181, State of California.
- Niehaus, T. 1977. Trifolium amoenum, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA
- Niehaus, T. and D. Fruchter. 1977. Navarretia plieantha, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA.
- Ornduff, R. 1954. Biosystematics of Blennosperma (Compositae). Brittonia 16, Vol. 16, 289-295.
- Ornduff, R. 1966. A biosystematic study of the goldfield genus Lasthenia. Univ. Calif. Publications in Botany, 40: 1 92.
- Ornduff, R. 1969. The origin and relationships of Lasthenia burkei (Compositae). American Journal of Botany 56 (9): 1042 1047.
- Ornduff, R. 1969. Limnanthes vinculans, a new California endemic. Brittonia 21: 11-14.
- Ornduff, R. 1977a. Lasthenia burkei, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA.

- Ornduff, R. 1977b. Blennosperma bakeri, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA.
- Patterson, C. A. 1987. Unpubl. report to Sonoma County Airport, botanical baseline inventory and mitigation plan for Burke's goldfields. Prep. for Sonoma Co. Airport.
- Patterson, C. A. 1990. Vernal pool creation and rare plant translocation in Sonoma County, California. Unpubl. report by C. Patterson, July 4, 1990.
- Pavlik, B., C. Muick, S. Johnson, and M. Popper. 1991. Oaks of California. Cachuma Press, Los Olivos, CA.
- Pennak, R. W. 1978. Freshwater invertebrates of the United States. Second edition; Wiley and Sons, New York. 803 pp.
- Raven, P. H., and D. I. Axelrod. 1978. Origin and relationships of the California flora. Univ. Calif. Publications in Botany, Vol. 72. Univ. Calif. Press, Berkeley. 134 pp.
- Remsen, J. V., Jr. 1978. Bird species of special concern. An annotated list of declining or vulnerable bird species. Prepared for Calif. Dept. Fish and Game (Sacramento) under project PR W-54-R-9 (Report No. 78-1), June, 1978.
- Resh, V. H. and D. M. Rosenberg (ed.s). 1984. The ecology of aquatic insects. Praeger Publishers, New York.
- Rubtzoff, P. 1977. Carex albida, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA.
- Rubtzoff, P. 1977. Rhynchospora californica, a status report.

 Unpubl. report prepared for the California Native Plant Society, Sacramento, CA.
- Santa Rosa, City of 1993. Southwest Area Plan. Specific area plan for southwest Santa Rosa; prepared 1993.
- Shaffer, H. B., R. N. Fisher, and S. E. Stanley. (no date Status Report: The California tiger salamander (Ambystoma californiense). Final report to the Calif. Dept. of Fish and Game, Inland fisheries Division Contracts FG 9422 and FG 1383.
- Skinner, M. W. and B. M. Pavlik. 1994. Inventory of rare and endangered vascular plants of California. California Native Plant Society Special Publication No. 1 (Fifth edition)
- Smith, J. P., Jr. and K. Berg. 1988. Inventory of rare and endangered vascular plants of California. California Native Plant Society Special Publication No. 1 (4th Edition)
- Soil Conservation Service. 1992. Hydric soils list: Santa Rosa Calif. Field Office. USDA-SCS, Davis, CA
- State of California. 1984. The California Environmental Quality Act.

 Laws and guidelines, January, 1984. Office of Planning and Research, State of California.
- Stebbins, G. L. and J. Major. 1965. Endemism and speciation in the California flora. Ecol. Monographs 35:1-35.

- Stebbins, R. C. 1985. A field guide to western reptiles and amphibians. A Peterson Field Guide. Houghton Mifflin Co., Boston
- Steinhart, P. 1990. California's wild heritage. Threatened and endangered animals in the golden state.

 Prepared for the California Dept. of Fish and Game, Calif. Academy of Sciences, and Sierra Club Books
- Thorne, R. F. 1981. Are California's vernal pools unique?In: Jain and Moyle (eds.). Vernal pools and intermittent streams. Univ. Calif. Institute of Ecology Publication No. 28
- Thorp, J. H. and A. P. Covich (ed.s). 1991. Ecology and classification of North American freshwater invertebrates. Academic Press, Inc. San Diego. 911 pp.
- U.S. Army Corps of Engineers. 1987. Corps of Engineers wetlands delineation Manual.
 Prep. by Environmental Laboratory, Dept. of the Army, Waterways Experiment Station; Wetlands Research Program, Tech. Report Y-87-1; Vicksburg, MI
- U.S. Fish and Wildlife Service. 1985. Endangered and threatened wildlife and plants; review of plant taxa for listing as endangered or threatened species; notice of review. 50 CFR Part 17. Federal Register Vol. 50, No. 188, Sept. 27, 1985.
- U.S. Fish and Wildlife Service. 1985. Wetland policy. Statement of regional FWS policy on wetlands, signed October 22, 1985. U.S. Fish and Wildlife Service, Sacramento, CA.
- U.S. Fish and Wildlife Service. 1989. Wetland plant list: Wetland plants of the State of California. Prep. by Porter B. Reed, Jr., Inland Freshwater Ecology Section, U.S. Fish and Wildlife Service, St. Petersburg, FL.
- U.S. Government. 1972. Federal water pollution control act amendments of 1972 (Public Law 92-500). Referred to as the Clean Water Act.
- Usinger, R. L. ed. 1956. Aquatic insects of California with keys to North American genera and California species. Univ. Calif. Press, Berkeley. 508 pp.
- Vandivere, W. 1993a. Hydrological analysis of the Bellevue School District site, Santa Rosa. Letter to Bellevue School District of July 20, 1993.
- Vandivere, W. 1993b. Hydrological analysis of the Wikiup Mitigation Bank site, Santa Rosa. Letter report to Wikiup Builders of July 12, 1993.
- Vandivere, W. 1994. Supplemental hydrological memo for the Wikiup mitigation Bank site, Santa Rosa. Letter report to Wikiup Builders of June, 1994.
- Waaland, M. E. 1989. Technical Memorandum No. W1: Baseline evaluation of Laguna de Santa Rosa wetlands and natural resources. Santa Rosa Subregional Water Reclamation System, long term detailed studies, prep. by CH2MHill.
- Waaland, M. E., M. Fawcett, J. Neilson, and D. W. Smith. 1990. Current condition and ecology of the Laguna ecosystem. In: Smith, D.W. (ed.), History, land uses, and natural resources of the Laguna de Santa Rosa. Prep. for the Santa Rosa Subregional Water Reclamation System.
- Waaland, M., R. M. Thompson, J. Vilms. 1989. Santa Rosa Plains endangered plant protection program report. Unpubl. report to Sonoma County Planning Dept. and Calif. Dept. of Fish and Game.